Canadian Journal of Anesthesia

Index of Authors

Volume 51, December 2004

Note: Abstracts and refresher course outlines in the June *Annual Meeting Supplement* are not included in this index.

A

Agarwal A, Dhiraj S, Raza M, Pandey R, Pandey CK, Singh PK, Singh U, Gupta D. Vein pretreatment with magnesium sulfate to prevent pain on injection of propofol is not justified, 130

Agarwal A. See *Pandey CK,* jt. auth.

Ahmad M. See *Munir MA,* jt. auth.

Akita M. See *Iida Y,* jt. auth.

Al-Refai A, Gunka V, Douglas J. Spinal anesthesia for Cesarean section in a parturient with long QT syndrome, 993

Albaladejo P. See *Geffroy A,* jt. auth.

Alhashemi JA, Khaki AM. Dexmedetomidine in combination with morphine PCA provides superior analgesia for shockwave lithotripsy, 342

Allaouchiche B. See *Bouvet L,* jt. auth.

Allard R, Hatzakorzian R, Deschamps A, Backman SB. Decreased heart rate and blood pressure in a recent cardiac transplant patient after spinal anesthesia, 829

Almuslim O, Butler R. Best evidence in critical care medicine: Selective digestive decontamination decreases mortality and morbidity in the intensive care. Commentary, 737

Almuslim O, Leasa D. Best evidence in critical care medicine: The use of recombinant human erythropoietin to reduce red cell transfusions in critically ill patients. Commentary, 621

Alvarez GG, Fergusson DA, Neilipovitz DT, Hébert PC. Cell salvage does not minimize perioperative allogeneic blood transfusion in abdominal vascular surgery: a systematic review, 425

Alvarez N. See *Dobson G,* jt. auth.

Ambesh SP. See *Pandey CK,* jt. auth.

Ansermino M. See *Minogue SC,* jt. auth.

Antonelli M. See *Conti G,* jt. auth.

Apfel CC. See *Biedler A,* jt. auth.; *Pierre S,* jt. auth.

Arcand G, Girard F, McCormack M, Chouinard P, Boudreault D, Williams S. Bilateral sixth cranial nerve palsy after unintentional dural puncture, 821

Arcangeli A. See *Conti G,* jt. auth.

Armstrong KPJ, Cherry RA. Brachial plexus anesthesia compared to general anesthesia when a block room is available, 41

Asai T, Shingu K. Airway management of a patient with tracheal stenosis for surgery in the prone position, 733

Atanassoff PG. See *Kurup V,* jt. auth.

Aubrun F, Hrazdilova O, Langeron O, Coriat P, Riou B. A high initial VAS score and sedation after *iv* morphine titration are associated with the need for rescue analgesia, 969

Avci A. See *Kanbak M,* jt. auth.

Aykaç B. See *Yüceyar L,* jt. auth.

Aypar U. See *Kanbak M,* jt. auth.

B

Babin D. See *Hemmerling TM,* jt. auth.

Backman SB. See *Allard R,* jt. auth.

Bahar M, Chanimov M, Cohen ML, Friedland M, Shul I, Gofman V, Gershfeld S, Geller R, Sherman DJ. The lateral recumbent head-down position decreases the incidence of epidural venous puncture during catheter insertion in obese parturients, 577

Balcioglu O. See *Gulev T,* jt. auth.

Balki M, Manninen PH. Craniotomy for suprasellar meningioma in a 28-week pregnant woman without fetal heart rate monitoring, 573

Barlow R, Karsli C, Luginbuehl I, Bissonnette B. Desflurane increases cerebral blood flow velocity when used for rapid emergence from propofol anesthesia in children, 824

Basile F. See *Hemmerling TM,* jt. auth.

Bau Y-J. See *Shin H-W,* jt. auth.

Baulig W. See *Dillier CM,* jt. auth.

Beaulieu P. See *Hemmerling TM,* jt. auth.

Beloeil H. See *Geffroy A,* jt. auth.

Benais H. See *Pierre S,* jt. auth.

Bendjelid K, Schutz N, Suter PM, Romand J-A. Après circulation extra-corporelle, la mesure continue de la Svo_2 et la co-oxymétrie du sang mêlé neonatal sont pas interchangeables / Continuous Svo_2 measurements and co-oximetry are not interchangeable immediately after cardiopulmonary bypass, 610

Benrhaiem M. See *Nathan N,* jt. auth.

Bérard J. See *Duflo F,* jt. auth.
Bernard F. See *Gorman T,* jt. auth.
Bernard N. See *Lopez S,* jt. auth.
Biedler A, Wermelt J, Kunitz O, Müller A, Wilhelm W, Dethling J, Apfel CC. A risk adapted approach reduces the overall institutional incidence of postoperative nausea and vomiting, 13
Bissonnette B. See *Barlow R,* jt. auth.
Blaine WC. See *Goldstein DH,* jt. auth.
Blanc C. See *Halkic N,* jt. auth.
Böhm SH. See *Tusman G,* jt. auth.
Boldt J. Fluid choice for resuscitation of the trauma patient: a review of the physiological, pharmacological, and clinical evidence, 500
Bonnaure-Mallet M. See *Tamanai-Shacoori Z,* jt. auth.
Bonnin P. See *Fattorutto M,* jt. auth.
Boselli E. See *Bouvet L,* jt. auth.*; Duflo F,* jt. auth.
Boudreault D. See *Arcand G,* jt. auth.*; Mac TB,* jt. auth.
Bouvet L, Allaouchiche B, Duflo F, Debon R, Chassard D, Boselli E. Le rémifentanil est une alternative efficace au propofol pour l'analgésie auto-contrôlée en endoscopie digestive / Remifentanil is an effective alternative to propofol for patient-controlled analgesia during digestive endoscopic procedures, 122
Bouvier E. See *Geffroy A,* jt. auth.
Bouvier G. See *Mac TB,* jt. auth.
Bowman KW, Richard SA. Cultural considerations for Canadians in the diagnosis of brain death, 273
Bozkurt K. See *Yüceyar L,* jt. auth.
Brill S, Plaza M. Non-narcotic adjuvants may improve the duration and quality of analgesia after knee arthroscopy: a brief review, 975
Brimacombe J, Stone T, Keller C. Supplementary cleaning does not remove protein deposits from re-usable laryngeal mask devices, 254
Brimacombe J. See *Keller C,* jt. auth.
Brouillette R. See *Koomson A,* jt. auth.
Brown KA. See *Koomson A,* jt. auth.
Brunetto GB. See *de Araujo DR,* jt. auth.
Bryson GL, Chung F, Cox RG, Crowe M-J, Fuller J, Henderson C, Finegan BA, Friedman Z, Miller DR, van Vlymen J. Patient selection in ambulatory anesthesia - An evidence-based review: part II, 782
Bryson GL, Chung F, Finegan BA, Friedman Z, Miller DR, van Vlymen J, Cox RG, Crowe M-J, Fuller J, Henderson C. Patient selection in ambulatory anesthesia - An evidence-based review: part I, 768
Buckley N. See *Yang H,* jt. auth.
Butler R. See *Almuslim O,* jt. auth.
Butterfield NN. See *Schwarz SKW,* jt. auth.

C

Campbell S. See *Kovacs G,* jt. auth.
Canadian Critical Care Trials Group. See *Sinuff T,* jt. auth.
Capdevila X. See *Lopez S,* jt. auth.
Cardinal V, Martin R, Tétrault J-P, Colas M-J, Gagnon L, Claprood Y. Bradycardie sévère et asystolie lors d'induction au sévoflurane et sufentanil à faible dose : à propos de trois cas / Low dose sufentanil and severe bradycardia during induction with sevoflurane: a report of three cases, 806
Carver ED. See *Crawford MW,* jt. auth.
Casey W. See *Iohom G,* jt. auth.
Castagnini HE, van Eijs F, Salevsky FC, Nathanson MH. Sevoflurane for interventional neuroradiology procedures is associated with more rapid early recovery than propofol, 486
Cavaliere F. See *Conti G,* jt. auth.
Cereda CMS. See *de Araujo DR,* jt. auth.
Chan RSN. See *Wong AYC,* jt. auth.
Chan W-H. See *Lin C-S,* jt. auth.
Chang S-H. See *Shin H-W,* jt. auth.
Chang Y-Y. See *Ting C-K,* jt. auth.
Chanimov M. See *Bahar M,* jt. auth.
Chari P. See *Mahajan R,* jt. auth.
Chassard D. See *Bouvet L,* jt. auth.*; Duflo F,* jt. auth.
Chaumeil A. See *Geffroy A,* jt. auth.
Chelly JE. See *Hagberg CA,* jt. auth.
Chen S. See *Zhang X,* jt. auth.
Cherry RA. See *Armstrong KPJ,* jt. auth.
Choi JH. See *Lee J,* jt. auth.
Choi PT-L. See *Yang H,* jt. auth.
Choinière J-L. See *Hemmerling TM,* jt. auth.
Chotel F. See *Duflo F,* jt. auth.
Chouinard P. See *Arcand G,* jt. auth.*; Mac TB,* jt. auth.
Chow L-H, Huang EY-K, Ho S-T, Lee T-Y, Tao P-L. Dextromethorphan potentiates morphine antinociception at the spinal level in rats, 905
Chua MV, Tsueda K, Doufas AG. Midazolam causes less sedation in volunteers with red hair, 25
Chua SMH, Sia ATH. Automated intermittent epidural boluses improve analgesia induced by intrathecal fentanyl during labour, 581
Chung DC. See *Ho AMH,* jt. auth.
Chung F, Tong D, Miceli PC, Reiz J, Harsanyi Z, Darke AC, Payne LW. Controlled-release codeine is equivalent to acetaminophen plus codeine for post-cholecystectomy analgesia, 216
Chung F. See *Bryson GL,* jt. auth.*; Friedman Z,* jt. auth.*; McGrath B,* jt. auth.
Claprood Y. See *Cardinal V,* jt. auth.
Cleave-Hogg D. See *Davies S,* jt. auth.

Cohen ML. See *Bahar M,* jt. auth.
Colas M-J. See *Cardinal V,* jt. auth.
Combes X, Dumerat M, Dhonneur G. Emergency gum elastic bougie-assisted tracheal intubation in four patients with upper airway distortion, 1022
Combet S. See *Duflo F,* jt. auth.
Conti G, Arcangeli A, Antonelli M, Cavaliere F, Costa R, Simeoni F, Proietti R. Sedation with sufentanil in patients receiving pressure support ventilation has no effects on respiration: a pilot study, 494
Cook D, Rocker G, Heyland D. Dying in the ICU: strategies that may improve end-of-life care, 266
Cook DJ. See *Rocker GM,* jt. auth.*; Sinuff T,* jt. auth.
Coriat P. See *Aubrun F,* jt. auth.
Corno G. See *Pierre S,* jt. auth.
Costa R. See *Conti G,* jt. auth.
Côté C. See *Kaloul I,* jt. auth.*; Lemay E,* jt. auth.
Cotter JT, Nielsen KC, Guller U, Steele SM, Klein SM, Greengrass RA, Pietrobon R. Increased body mass index and ASA physical status IV are risk factors for block failure in ambulatory surgery - an analysis of 9,342 blocks, 810
Coulic V. See *Guntz E,* jt. auth.
Couture P. See *Denault A,* jt. auth.
Cox RG. Repair of incarcerated inguinal hernia in an infant with acute viral bronchiolitis, 68
Cox RG. See *Bryson GL,* jt. auth.
Crawford MW, Speakman M, Carver ED, Kim PCW. Acute chest syndrome shows a predilection for basal lung regions on the side of upper abdominal surgery, 707
Creinin MD. See *Ray BK,* jt. auth.
Crowe M-J. See *Bryson GL,* jt. auth.
Crystal GJ, Metwally AA, Salem MR. Isoflurane preserves central nervous system blood flow during intraoperative cardiac tamponade in dogs, 1011
Cupa M. See *Fattorutto M,* jt. auth.
Curti B. See *McGrath B,* jt. auth.

D

Dagenais P. See *Gorman T,* jt. auth.
Daper A. See *Van der Linden PJ,* jt. auth.
Darke AC. See *Chung F,* jt. auth.
Davies S, Cleave-Hogg D. Continuing medical education should be offered by both e-mail and regular mail: a survey of Ontario anesthesiologists, 444
de Araujo DR, Cereda CMS, Brunetto GB, Pinto LMA, Santana MHA, de Paula E. Encapsulation of mepivacaine prolongs the analgesia provided by sciatic nerve blockade in mice, 566
De Hert SG. See *Van der Linden PJ,* jt. auth.
de Moerloose P. See *Hardy J-F,* jt. auth.
de Paula E. See *de Araujo DR,* jt. auth.
Debizet E. See *Guntz E,* jt. auth.
Debon R. See *Bouvet L,* jt. auth.
Defrance P. See *Van der Linden PJ,* jt. auth.
Deibl M. See *Smolle M,* jt. auth.
Delbos A. See *Guntz E,* jt. auth.
Delhaye D. See *Guntz E,* jt. auth.
Denault A, Couture P, Pellerin M. 3D systolic anterior motion of the mitral valve (Images in anesthesia), 481
Denault AY. See *Williams SR,* jt. auth.
Deschamps A. See *Allard R,* jt. auth.
Deschamps S. See *Hemmerling TM,* jt. auth.
Dethling J. See *Biedler A,* jt. auth.
Devereaux PJ, Leslie K. Best evidence in anesthetic practice: Prevention: α- and β-adrenergic antagonists reduce perioperative cardiac events. Commentary, 290
Dhamee MS. See *Yao L,* jt. auth.
Dhiraaj S. See *Pandey CK,* jt. auth.
Dhiraj S. See *Agarwal A,* jt. auth.
Dhonneur G. See *Combes X,* jt. auth.
Dilger JA, Sprung J, Maurer W, Tetzlaff J. Remifentanil provides better analgesia than alfentanil during breast biopsy surgery under monitored anesthesia care, 20
Dillier CM, Trachsel D, Baulig W, Gysin C, Gerber AC, Weiss M. Laryngeal damage due to an unexpectedly large and inappropriately designed cuffed pediatric tracheal tube in a 13-month-old child, 72
Dobson G, Petrasek P, Alvarez N. Transesophageal echocardiography enhances endovascular stent placement in traumatic trans-section of the thoracic aorta (Images in anesthesia), 931
Dodek PM. See *Rocker GM,* jt. auth.*; Sinuff T,* jt. auth.
Dohi S. See *Iizawa A,* jt. auth.
Donati F. See *Hemmerling TM,* jt. auth.
Doufas AG. See *Chua MV,* jt. auth.
Douglas J. See *Al-Refai A,* jt. auth.
Douville C. See *Lee LA,* jt. auth.
Drolet P. See *Turcotte C,* jt. auth.
Duflo F, Qamouss Y, Rémond C, Pouyau A, Heilporn A, Taylor P, Paturel B, Combet S, Boselli E, Chotel F, Bérard J, Chassard D. Patient-controlled regional analgesia is effective in children: a preliminary report, 928
Duflo F. See *Bouvet L,* jt. auth.
Dugas G, Fuller J, Singh S, Watson J. Pheochromocytoma and pregnancy: a case report and review of anesthetic management, 134
Duggan S. See *Rodriguez RA,* jt. auth.

Dullenkopf A, Gerber A, Weiss M. The Microcuff™ tube allows a longer time interval until unsafe cuff pressures are reached in children, 997
Dumerat M. See *Combes X,* jt. auth.

E
Easton DM. See *Olafson K,* jt. auth.
El-Moalem HE. See *Habib AS,* jt. auth.
Elgendy H. See *McGrath B,* jt. auth.
Erolçay H. See *Yüceyar L,* jt. auth.
Eustace NP. See *Harmon DC,* jt. auth.

F
Fallaha M. See *Kaloul I,* jt. auth.
Fattorutto M, Tourreau-Pham S, Mazoyer E, Bonnin P, Raphaël M, Morin F, Cupa M, Samama C-M. Recombinant activated factor VII decreases bleeding without increasing arterial thrombosis in rabbits, 672
Feiss P. See *Nathan N,* jt. auth.
Fergusson D, Khanna MP, Tinmouth A, Hébert PC. Transfusion of leukoreduced red blood cells may decrease postoperative infections: two meta-analyses of randomized controlled trials, 417
Fergusson DA. See *Alvarez GG,* jt. auth.
Fick GH. See *Maltby JR,* jt. auth.
Finegan BA. See *Bryson GL,* jt. auth.
Finucane B. See *Tsui BCH,* jt. auth.
Fisher MM. See *Sinuff T,* jt. auth.
Fortier JD. See *Hemmerling TM,* jt. auth.
Franciosi LG. See *Schwarz SKW,* jt. auth.
Franklin R. See *Iohom G,* jt. auth.
Freud E. See *Shenkman Z,* jt. auth.
Friedland M. See *Bahar M,* jt. auth.
Friedman Z, Chung F, Wong DT. Ambulatory surgery adult patient selection criteria - a survey of Canadian anesthesiologists, 437
Friedman Z. See *Bryson GL,* jt. auth.
Fuller J. See *Bryson GL,* jt. auth.*; Dugas G,* jt. auth.
Furuya H. See *Matsuda T,* jt. auth.

G
Gaber O. See *Hinteregger M,* jt. auth.
Gagnon L. See *Cardinal V,* jt. auth.
Gan TJ. See *Habib AS,* jt. auth.
Geffroy A, Beloeil H, Bouvier E, Chaumeil A, Albaladejo P, Marty J. Prolonged right ventricular failure after relief of cardiac tamponade, 482
Geller R. See *Bahar M,* jt. auth.
Gerber A. See *Dullenkopf A,* jt. auth.
Gerber AC. See *Dillier CM,* jt. auth.
Gershfeld S. See *Bahar M,* jt. auth.
Ghori KG. See *Harmon DC,* jt. auth.
Gillet M. See *Halkic N,* jt. auth.
Girard F. See *Arcand G,* jt. auth.*; Mac TB,* jt. auth.
Girard M. See *Turcotte C,* jt. auth.
Goeschl A. See *Hinteregger M,* jt. auth.
Gofman V. See *Bahar M,* jt. auth.
Goldstein DH, VanDenKerkhof EG, Blaine WC. Acute pain management services have progressed, albeit insufficiently in Canadian academic hospitals, 231
Goldstein DH. See *VanDenKerkhof EG,* jt. auth.
Gorman T, Bernard F, Marquis F, Dagenais P, Skrobik Y. Best evidence in critical care medicine: Daily interruption of sedative infusions in critically ill patients undergoing mechanical ventilation. Commentary, 492
Greengrass RA. See *Cotter JT,* jt. auth.
Griffith LE. See *Sinuff T,* jt. auth.
Groupe d'Intérêt en Hémostase Périopératoire. See *Hardy J-F,* jt. auth.
Grover VK. See *Mahajan R,* jt. auth.
Guay J, Lortie L. An evaluation of pediatric in-hospital advanced life support interventions using the pediatric Utstein guidelines: a review of 203 cardiorespiratory arrests, 373
Guay J. See *Kaloul I,* jt. auth.*; Lemay E,* jt. auth.
Gulev T, Unlugenc H, Gundogan Z, Ozalevli M, Balcioglu O, Topcuoglu MS. A background infusion of morphine enhances patient-controlled analgesia after cardiac surgery, 718
Guller U. See *Cotter JT,* jt. auth.
Gundogan Z. See *Gulev T,* jt. auth.
Gunka V. See *Al-Refai A,* jt. auth.
Guntz E, Herman P, Debizet E, Delhaye D, Coulic V, Sosnowski M. Sciatic nerve block in the popliteal fossa: description of a new medial approach, 817
Guntz E, Herman P, Delbos A, Sosnowski M. The radial nerve should be blocked before the ulnar nerve during a brachial plexus block at the humeral canal, 354
Gupta D. See *Agarwal A,* jt. auth.*; Pandey CK,* jt. auth.
Guyatt GH. See *Sinuff T,* jt. auth.
Gysin C. See *Dillier CM,* jt. auth.

H
Ha DV, Johnson D. High frequency oscillatory ventilation in the management of a high output bronchopleural fistula: a case report, 78
Habib AS, El-Moalem HE, Gan TJ. The efficacy of the 5-HT_3 receptor antagonists combined with droperidol for PONV prophylaxis is similar to their combination with dexamethasone. A meta-analysis of randomized controlled trials, 311

Habib AS, Gan TJ. Evidence-based management of postoperative nausea and vomiting: a review, 326
Hagberg CA, Vartazarian TN, Chelly JE, Ovassapian A. The incidence of gastroesophageal reflux and tracheal aspiration detected with pH electrodes is similar with the Laryngeal Mask Airway® and Esophageal Tracheal Combitube® - a pilot study, 243
Halkic N, Ksontini R, Scholl B, Blanc C, Kovacsovics T, Meylan P, Muheim C, Gillet M, Mosimann F. Recurrent cytomegalovirus disease, visceral leishmaniosis, and Legionella pneumonia after liver transplantation: a case report, 84
Hall LE. See *Rodriguez RA,* jt. auth.
Hall RI, Rocker GM, Murray D. Simple changes can improve conduct of end-of-life care in the intensive care unit, 631
Hall SRR, Wang L, Milne B, Hong M. Central dexmedetomidine attenuates cardiac dysfunction in a rodent model of intracranial hypertension, 1025
Halwagi A. See *Kaloul I,* jt. auth.
Hanaoka K. See *Nishiyama T,* jt. auth.
Hardy J-F, de Moerloose P, Samama M, Members of the Groupe d'Intérêt en Hémostase Périopératoire. Massive transfusion and coagulopathy: pathophysiology and implications for clinical management, 293
Harmon DC, Ghori KG, Eustace NP, O'Callaghan SJF, O'Donnell AP, Shorten GD. Aprotinin decreases the incidence of cognitive deficit following CABG and cardiopulmonary bypass: a pilot randomized controlled study, 1002
Harsanyi Z. See *Chung F,* jt. auth.
Hastings RH. See *Lee J,* jt. auth.
Hatano Y. See *Yu J,* jt. auth.
Hatzakorzian R. See *Allard R,* jt. auth.
Hébert PC. See *Alvarez GG,* jt. auth.*; Fergusson D,* jt. auth.
Heilporn A. See *Duflo F,* jt. auth.
Hemmerling TM, Beaulieu P, Jacobi KE, Babin D, Schmidt J. Neuromuscular blockade does not change the incidence or severity of pharyngolaryngeal discomfort after LMA anesthesia, 728
Hemmerling TM, Michaud G, Babin D, Trager G, Donati F. Comparison of phonomyography with balloon pressure mechanomyography to measure contractile force at the corrugator supercilii muscle, 116
Hemmerling TM, Michaud G, Trager G, Deschamps S. Phonomyographic measurements of neuromuscular blockade are similar to mechanomyography for hand muscles, 795
Hemmerling TM, Prieto I, Choinière J-L, Basile F, Fortier JD. Ultra-fast-track anesthesia in off-pump coronary artery bypass grafting: a prospective audit comparing opioid-based anesthesia *vs* thoracic epidural-based anesthesia, 163
Henderson C. See *Bryson GL,* jt. auth.
Herman P. See *Guntz E,* jt. auth.
Heyland D. See *Cook D,* jt. auth.
Heyland DK. See *Rocker GM,* jt. auth.
Hinteregger M, Zschiegner F, Lirk P, Ladner E, Goeschl A, Gaber O, Moser P, Lorenz I, Kolbitsch C. A new guidance device facilitates percutaneous puncture of the foramen ovale in human cadavers, 990
Ho AMH, Chung DC, To EWH, Karmakar MK. Total airway obstruction during local anesthesia in a non-sedated patient with a compromised airway, 838
Ho S-T. See *Chow L-H,* jt. auth.
Hong M. See *Hall SRR,* jt. auth.
Hoppenstein D. See *Shenkman Z,* jt. auth.
Horishita T. See *Minami K,* jt. auth.*; Ogata J,* jt. auth.
Hrazdilova O. See *Aubrun F,* jt. auth.
Hu JS. See *Ting C-K,* jt. auth.
Hu W. See *Zhang X,* jt. auth.
Huang EY-K. See *Chow L-H,* jt. auth.

I

Iida Y, Akita M, Nukariya M. Temporary atrial pacing via a triple lumen central venous catheter (Images in anesthesia), 242
Iijima T, Takagi T, Iwao Y. An increased circulating blood volume does not prevent hypotension after pheochromocytoma resection, 212
Iizawa A, Oshima T, Kasuya Y, Dohi S. Oral tandospirone and clonidine provide similar relief of preoperative anxiety, 668
Inoue S. See *Matsuda T,* jt. auth.
Iohom G, Franklin R, Casey W, Lyons B. The McCoy straight blade does not improve laryngoscopy and intubation in normal infants, 155
Irwin MG. See *Wong AYC,* jt. auth.
Ishida S. See *Sasano N,* jt. auth.
Ishikawa K. See *Sasano N,* jt. auth.
Iwahashi S. See *Yu J,* jt. auth.
Iwao Y. See *Iijima T,* jt. auth.

J

Jacobi KE. See *Hemmerling TM,* jt. auth.
Jedeikin R. See *Shenkman Z,* jt. auth.
Johnson D. See *Ha DV,* jt. auth.
Jones G. See *Meggison H,* jt. auth.
Joshi M. See *Manninen P,* jt. auth.

K

Kaloul I, Guay J, Côté C, Fallaha M. The posterior lumbar plexus (psoas compartment) block and the three-in-one femoral nerve block provide similar postoperative analgesia after total knee replacement, 45

Kaloul I, Guay J, Côté C, Halwagi A, Varin F. Ropivacaine plasma concentrations are similar during continuous lumbar plexus blockade using the anterior three-in-one and the posterior psoas compartment techniques, 52

Kalow W. Atypical plasma cholinesterase. A personal discovery story: a tale of three cities, 206

Kamming D. See *McGrath B,* jt. auth.

Kanaya N. See *Sugino S,* jt. auth.

Kanazawa T. See *Takeda S,* jt. auth.

Kanbak M, Saricaoglu F, Avci A, Ocal T, Koray Z, Aypar U. Propofol offers no advantage over isoflurane anesthesia for cerebral protection during cardiopulmonary bypass: a preliminary study of S-100β protein levels, 712

Karmakar MK. See *Ho AMH,* jt. auth.

Karsli C. See *Barlow R,* jt. auth.

Kasuya Y. See *Iizawa A,* jt. auth.

Katsuya H. See *Sasano N,* jt. auth.

Kaul B. See *Ray BK,* jt. auth.

Keiichi S. See *Matsuda T,* jt. auth.

Keller C, Brimacombe J, Moriggl B, Lirk P, von Goedeck A. In cadavers, directly measured mucosal pressures are similar for the Unique™ and the Soft Seal™ laryngeal mask airway devices, 834

Keller C. See *Brimacombe J,* jt. auth.*; Smolle M,* jt. auth.

Khaki AM. See *Alhashemi JA,* jt. auth.

Khanna MP. See *Fergusson D,* jt. auth.

Kim ES. See *Lee J,* jt. auth.

Kim EY. See *Schwarz SKW,* jt. auth.

Kim PCW. See *Crawford MW,* jt. auth.

Kimbimbi P. See *Van der Linden PJ,* jt. auth.

King S. See *McGrath B,* jt. auth.

Kirk P. See *Ramachandran K,* jt. auth.

Klein SM. See *Cotter JT,* jt. auth.

Klemenzson GK, Perouansky M. Contemporary anesthesia ventilators incur a significant "oxygen cost," 616

Kolbitsch C. See *Hinteregger M,* jt. auth.

Konen O. See *Shenkman Z,* jt. auth.

Konukoglu D. See *Yüceyar L,* jt. auth.

Koomson A, Morin I, Brouillette R, Brown KA. Children with severe OSAS who have adenotonsillectomy in the morning are less likely to have postoperative desaturation than those operated in the afternoon, 62

Koray Z. See *Kanbak M,* jt. auth.

Kovacs G, Law JA, Ross J, Tallon J, MacQuarrie K, Petrie D, Campbell S, Soder C. Acute airway management in the emergency department by non-anesthesiologists, 174

Kovacsovics T. See *Halkic N,* jt. auth.

Kraus PA. See *Sinuff T,* jt. auth.

Ksontini R. See *Halkic N,* jt. auth.

Kuczkowski KM. The cocaine abusing parturient: a review of anesthetic considerations, 145

Kunitz O. See *Biedler A,* jt. auth.

Kurup V, Ramani R, Atanassoff PG. Sedation after spinal anesthesia in elderly patients: a preliminary observational study with the PSA-4000, 562

Kutsogiannis DJ. See *Rocker GM,* jt. auth.

Kwon OK. See *Lee J,* jt. auth.

L

Ladner E. See *Hinteregger M,* jt. auth.

Lam AM. See *Lee LA,* jt. auth.

Langeron O. See *Aubrun F,* jt. auth.

Law JA. See *Kovacs G,* jt. auth.

Lazar NM. See *Sinuff T,* jt. auth.

Leasa D. See *Almuslim O,* jt. auth.

Lee H-W. See *Shin H-W,* jt. auth.

Lee J, Choi JH, Lee YK, Kim ES, Kwon OK, Hastings RH. The Callander laryngoscope blade modification is associated with a decreased risk of dental contact, 181

Lee LA, Vavilala MS, Lam AM, Douville C, Mooret A, Visco E, Newell DW. Ophthalmic artery blood flow velocity increases during hypocapnia, 388

Lee T-Y. See *Chow L-H,* jt. auth.

Lee YK. See *Lee J,* jt. auth.

Lemay E, Guay J, Côté C, Roy A. Tranexamic acid reduces the need for allogenic red blood cell transfusions in patients undergoing total hip replacement, 31

Leslie K. See *Devereaux PJ,* jt. auth.

Level of Care Study Investigators. See *Sinuff T,* jt. auth.

Levy MM. See *Sinuff T,* jt. auth.

Lim H-J. See *Shin H-W,* jt. auth.

Lin C-J. See *Lin C-S,* jt. auth.

Lin C-S, Sun W-Z, Chan W-H, Lin C-J, Yeh H-M, Mok MS. Intravenous lidocaine and ephedrine, but not propofol, suppress fentanyl-induced cough, 654

Lirk P. See *Hinteregger M,* jt. auth.*; Keller C,* jt. auth.*; Smolle M,* jt. auth.

Littleford J. Effects on the fetus and newborn of maternal analgesia and anesthesia: a review, 586

Liu Y. See *Zhang X,* jt. auth.

Lopez S, Privat A, Bernard N, Ohanna F, Vergnes C, Capdevila X. Intrathecal bupivacaine protects against extension of lesions in an acute photochemical spinal cord injury model, 364
Lorenz I. See *Hinteregger M,* jt. auth.
Lortie L. See *Guay J,* jt. auth.
Luginbuehl I. See *Barlow R,* jt. auth.
Lyons B. See *Iohom G,* jt. auth.

M
Mac TB, Girard F, McKenty S, Chouinard P, Boudreault D, Ruel M, Bouvier G. A difficult airway is not more prevalent in patients suffering from spasmodic torticollis: a case series, 250
MacLeod BA. See *Schwarz SKW,* jt. auth.
MacQuarrie K. See *Kovacs G,* jt. auth.
Mahajan R, Grover VK, Chari P. Caudal neostigmine with bupivacaine produces a dose-independent analgesic effect in children, 702
Maltby JR, Pytka S, Watson NC, McTaggart Cowan RA, Fick GH. Drinking 300 mL of clear fluid two hours before surgery has no effect on gastric fluid volume and pH in fasting and non-fasting obese patients, 111
Mandell GL. See *Ray BK,* jt. auth.
Manninen P, Sarjeant R, Joshi M. Posterior tibial nerve and median nerve somatosensory evoked potential monitoring during carotid endarterectomy, 937
Manninen PH. See *Balki M,* jt. auth.
Marquet P. See *Nathan N,* jt. auth.
Marquis F. See *Gorman T,* jt. auth.
Marshall JC. See *Sinuff T,* jt. auth.
Martin R. See *Cardinal V,* jt. auth.
Martineau R. See *Williams SR,* jt. auth.
Martinek RM. Witnessed asystole during spinal anesthesia treated with atropine and ondansetron: a case report, 226
Marty J. See *Geffroy A,* jt. auth.
Masuda R. See *Takeda S,* jt. auth.
Matioc AA, Olson J. Use of the Laryngeal Tube™ in two unexpected difficult airway situations: lingual tonsillar hyperplasia and morbid obesity, 1018
Matsuda T, Inoue S, Keiichi S, Furuya H. Accidental knot formation of a pulmonary artery catheter (Images in anesthesia), 1010
Matsukawa T. See *Nishiyama T,* jt. auth.
Maurer W. See *Dilger JA,* jt. auth.
Mazoyer E. See *Fattorutto M,* jt. auth.
McChesney J. See *Yang H,* jt. auth.
McCormack M. See *Arcand G,* jt. auth.
McDermid RC. Best evidence in critical care medicine: Treatment of submassive pulmonary embolism. Commentary, 846
McDonald E. See *Sinuff T,* jt. auth.
McDonald WN. See *Schwarz SKW,* jt. auth.
McGrath B, Elgendy H, Chung F, Kamming D, Curti B, King S. Thirty percent of patients have moderate to severe pain 24 hr after ambulatory surgery: a survey of 5,703 patients, 886
McKenty S. See *Mac TB,* jt. auth.
McTaggart Cowan RA. See *Maltby JR,* jt. auth.
Meggison H, Jones G. Best evidence in critical care medicine: Treatment: adrenal replacement therapy improves survival in patients with septic shock. Commentary, 264
Metwally AA. See *Crystal GJ,* jt. auth.
Meylan P. See *Halkic N,* jt. auth.
Miceli PC. See *Chung F,* jt. auth.
Michaud G. See *Hemmerling TM,* jt. auth.
Miller DR. See *Bryson GL,* jt. auth.
Milne B. See *Hall SRR,* jt. auth.*; VanDenKerkhof EG,* jt. auth.
Minami K, Ogata J, Horishita T, Shiraishi M, Okamoto T, Sata T, Shigematsu A. Intramuscular tramadol increases gastric pH during anesthesia, 545
Minami K. See *Ogata J,* jt. auth.
Minogue SC, Morrisson M, Ansermino M. Laryngo-tracheo-bronchial stenosis in a patient with primary pulmonary amyloidosis: a case report and brief review, 842
Missant C, Teunkens A, Vandermeersch E, Van de Velde M. Intrathecal clonidine prolongs labour analgesia but worsens fetal outcome: a pilot study, 696
Miyamoto H. See *Ogata J,* jt. auth.
Mizuuchi M. See *Sugino S,* jt. auth.
Mok MS. See *Lin C-S,* jt. auth.
Mondragon SA. See *Waxler B,* jt. auth.
Mooret A. See *Lee LA,* jt. auth.
Moriggl B. See *Keller C,* jt. auth.
Morin F. See *Fattorutto M,* jt. auth.
Morin I. See *Koomson A,* jt. auth.
Morrisson M. See *Minogue SC,* jt. auth.
Moser P. See *Hinteregger M,* jt. auth.
Mosimann F. See *Halkic N,* jt. auth.
Muheim C. See *Halkic N,* jt. auth.
Müller A. See *Biedler A,* jt. auth.
Munir MA, Zhang J, Ahmad M. A modified needle-inside-needle technique for the ganglion impar block, 915
Murray D. See *Hall RI,* jt. auth.

N

Nakayama M. See *Sugino S,* jt. auth.

Namiki A. See *Sugino S,* jt. auth.

Nathan N, Vandroux D, Benrhaiem M, Marquet P, Preux P-M, Feiss P. Low alfentanil target-concentrations improve hemodynamic and intubating conditions during induction with sevoflurane, 382

Nathanson MH. See *Castagnini HE,* jt. auth.

Nedumgottil K. See *Waxler B,* jt. auth.

Neilipovitz DT. See *Alvarez GG,* jt. auth.

Newell DW. See *Lee LA,* jt. auth.

Nielsen KC. See *Cotter JT,* jt. auth.

Nishiyama T, Hanaoka K. The A-line ARX index may be a more sensitive detector of arousal than the bispectral index during propofol-fentanyl-nitrous oxide anesthesia: a preliminary investigation, 539

Nishiyama T, Matsukawa T, Hanaoka K. Intrathecal propofol has analgesic effects on inflammation-induced pain in rats, 899

Nukariya M. See *Iida Y,* jt. auth.

O

Ocal T. See *Kanbak M,* jt. auth.

O'Callaghan CJ. See *Rocker GM,* jt. auth.

O'Callaghan SJF. See *Harmon DC,* jt. auth.

Ocampo CE. See *Supandji M,* jt. auth.

O'Donnell AP. See *Harmon DC,* jt. auth.

Ogata J, Minami K, Miyamoto H, Horishita T, Ogawa M, Sata T, Taniguchi H. Gargling with povidone-iodine reduces the transport of bacteria during oral intubation, 932

Ogata J. See *Minami K,* jt. auth.

Ogawa K. See *Yu J,* jt. auth.

Ogawa M. See *Ogata J,* jt. auth.

Ohanna F. See *Lopez S,* jt. auth.

Okamoto T. See *Minami K,* jt. auth.

Olafson K, Selaman M, Easton DW. BECCM Best evidence in critical care medicine: Therapeutic hypothermia to improve neurologic outcome after cardiac arrest. Commentary, 76

Olson J. See *Matioc AA,* jt. auth.

Oshima T. See *Iizawa A,* jt. auth.

Ovassapian A. See *Hagberg CA,* jt. auth.

Ozalevli M. See *Gulev T,* jt. auth.

P

Pandey CK, Priye S, Singh S, Singh U, Singh RB, Singh PK. Preemptive use of gabapentin significantly decreases postoperative pain and rescue analgesic requirements in laparoscopic cholecystectomy, 358

Pandey CK, Raza M, Dhiraaj S, Agarwal A, Singh PK. Rapid preparation of severe uncontrolled thyrotoxicosis due to Graves' disease with Iopanoic acid - case report, 38

Pandey CK, Sahay S, Gupta D, Ambesh SP, Singh RB, Raza M, Singh U, Singh PK. Preemptive gabapentin decreases postoperative pain after lumbar discoidectomy, 986

Pandey CK. See *Agarwal A,* jt. auth.

Pandey R. See *Agarwal A,* jt. auth.

Parlow JL. See *VanDenKerkhof EG,* jt. auth.

Parry G. Trendelenburg position, head elevation and a midline position optimize right internal jugular vein diameter, 379

Patel SN. See *Waxler B,* jt. auth.

Paturel B. See *Duflo F,* jt. auth.

Payne LW. See *Chung F,* jt. auth.

Pellerin M. See *Denault A,* jt. auth.*; Williams SR,* jt. auth.

Perouansky M. See *Klemenzson GK,* jt. auth.

Petrasek P. See *Dobson G,* jt. auth.

Petrie D. See *Kovacs G,* jt. auth.

Pierre S, Corno G, Benais H, Apfel CC. A risk score-dependent antiemetic approach effectively reduces postoperative nausea and vomiting - a continuous quality improvement initiative, 320

Pietrobon R. See *Cotter JT,* jt. auth.

Pinggera G. See *Smolle M,* jt. auth.

Pinto LMA. See *de Araujo DR,* jt. auth.

Plaza M. See *Brill S,* jt. auth.

Pouyau A. See *Duflo F,* jt. auth.

Preux P-M. See *Nathan N,* jt. auth.

Prieto I. See *Hemmerling TM,* jt. auth.

Privat A. See *Lopez S,* jt. auth.

Priye S. See *Pandey CK,* jt. auth.

Proietti R. See *Conti G,* jt. auth.

Pytka S. See *Maltby JR,* jt. auth.

Q

Qamouss Y. See *Duflo F,* jt. auth.

R

Rabuka S. See *Rapchuk I,* jt. auth.

Ramachandran K, Kirk P. Massive hemorrhage in a previously undiagnosed abdominal pregnancy presenting for elective Cesarean delivery, 57

Ramanathan S. See *Ray BK,* jt. auth.

Ramani R. See *Kurup V,* jt. auth.

Rapchuk I, Rabuka S, Tonelli M. Perioperative use of beta-blockers remains low: experience of a single Canadian tertiary institution, 761

Raphaël M. See *Fattorutto M,* jt. auth.

Rathaus V. See *Shenkman Z,* jt. auth.

Ray BK, Vallejo MC, Creinin MD, Shannon KT, Mandell GL, Kaul B, Ramanathan S. Amniotic fluid embolism with second trimester pregnancy termination: a case report, 139
Raza M, Dhiraaj S, Agarwal A, Singh PK. Rapid preparation of severe uncontrolled thyrotoxicosis due to Graves' disease with Iopanoic acid - case report, 38
Raza M. See *Agarwal A,* jt. auth.*; Pandey CK,* jt. auth.
Reiz J. See *Chung F,* jt. auth.
Rémond C. See *Duflo F,* jt. auth.
Richard SA. See *Bowman KW,* jt. auth.
Rieder J. See *Smolle M,* jt. auth.
Ries CR. See *Schwarz SKW,* jt. auth.
Riou B. See *Aubrun F,* jt. auth.
Robert J-C. See *Tamanai-Shacoori Z,* jt. auth.
Rochon AG. See *Tessler MJ,* jt. auth.
Rocker G. See *Cook D,* jt. auth.
Rocker GM, Heyland DK, Cook DJ, Dodek PM, Kutsogiannis DJ, O'Callaghan CJ. Most critically ill patients are perceived to die in comfort during withdrawal of life support: a Canadian multicentre study, 623
Rocker GM. See *Hall RI,* jt. auth.*; Sinuff T,* jt. auth.
Rodriguez RA, Hall LE, Duggan S, Splinter WM. The bispectral index does not correlate with clinical signs of inhalational anesthesia during sevoflurane induction and arousal in children, 472
Romand J-A. See *Bendjelid K,* jt. auth.
Ross J. See *Kovacs G,* jt. auth.
Roy A. See *Lemay E,* jt. auth.
Ruel M. See *Mac TB,* jt. auth.

S

Sahay S. See *Pandey CK,* jt. auth.
Salem MR. See *Crystal GJ,* jt. auth.
Salevsky FC. See *Castagnini HE,* jt. auth.
Samama C-M. See *Fattorutto M,* jt. auth.
Samama M. See *Hardy J-F,* jt. auth.
Santana MHA. See *de Araujo DR,* jt. auth.
Saricaoglu F. See *Kanbak M,* jt. auth.
Sarjeant R. See *Manninen P,* jt. auth.
Sasano H. See *Sasano N,* jt. auth.
Sasano N, Ishida S, Tetsu S, Takasu H, Ishikawa K, Sasano H, Katsuya H. Cerebral fat embolism diagnosed by magnetic resonance imaging at one, eight, and 50 days after hip arthroplasty: a case report, 875
Sata T. See *Minami K,* jt. auth.*; Ogata J,* jt. auth.
Schmartz D. See *Van der Linden PJ,* jt. auth.
Schmidt J. See *Hemmerling TM,* jt. auth.
Scholl B. See *Halkic N,* jt. auth.
Schutz N. See *Bendjelid K,* jt. auth.
Schwarz SKW, Butterfield NN, MacLeod BA, Kim EY, Franciosi LG, Ries CR. Under "real world" conditions, desflurane increases drug cost without speeding discharge after short ambulatory anesthesia compared to isoflurane, 892
Schwarz SKW, Wong CL, McDonald WN. Spontaneous recovery from a spinal epidural hematoma with atypical presentation in a nonagenarian, 557
Selaman M. See *Olafson K,* jt. auth.
Shacoori V. See *Tamanai-Shacoori Z,* jt. auth.
Shannon KT. See *Ray BK,* jt. auth.
Shenkman Z, Rathaus V, Jedeikin R, Konen O, Hoppenstein D, Snyder M, Freud E. The distance from the skin to the subarachnoid space can be predicted in premature and former-premature infants, 160
Sherman DJ. See *Bahar M,* jt. auth.
Shigematsu A. See *Minami K,* jt. auth.
Shin H-W, Bau Y-J, Lee H-W, Lim H-J, Yoon S-M, Chang S-H. Arousal with *iv* epinephrine depends on the depth of anesthesia, 880
Shingu K. See *Asai T,* jt. auth.
Shiraishi M. See *Minami K,* jt. auth.
Shorten GD. See *Harmon DC,* jt. auth.
Shrier I. See *Tessler MJ,* jt. auth.
Shul I. See *Bahar M,* jt. auth.
Sia ATH. See *Chua SMH,* jt. auth.*; Supandji M,* jt. auth.
Simeoni F. See *Conti G,* jt. auth.
Singh PK. See *Agarwal A,* jt. auth.*; Pandey CK,* jt. auth.
Singh RB. See *Pandey CK,* jt. auth.
Singh S. See *Dugas G,* jt. auth.*; Pandey CK,* jt. auth.
Singh U. See *Agarwal A,* jt. auth.*; Pandey CK,* jt. auth.
Sinuff T, Cook DJ, Rocker GM, Griffith LE, Walter SD, Fisher MM, Dodek PM, Sjokvist P, McDonald E, Marshall JC, Kraus PA, Levy MM, Lazar NM, Guyatt GH, Level of Care Study Investigators, Canadian Critical Care Trials Group. DNR directives are established early in mechanically ventilated intensive care unit patients, 1034
Sjokvist P. See *Sinuff T,* jt. auth.
Skjodt NM, Townsend DR. Best evidence in critical care medicine: Treatment: vasodilatation with nitroprusside in symptomatic patients with severe aortic stenosis and congestive heart failure improves cardiac index. Commentary, 1042
Skrobik Y. See *Gorman T,* jt. auth.

Smolle M, Keller C, Pinggera G, Deibl M, Rieder J, Lirk P. Clear hydro-gel, compared to ointment, provides improved eye comfort after brief surgery, 126
Snyder M. See *Shenkman Z,* jt. auth.
Soder C. See *Kovacs G,* jt. auth.
Sosnowski M. See *Guntz E,* jt. auth.
Speakman M. See *Crawford MW,* jt. auth.
Splinter WM. See *Rodriguez RA,* jt. auth.
Sprung J. See *Dilger JA,* jt. auth.
Steele SM. See *Cotter JT,* jt. auth.
Stone T. See *Brimacombe J,* jt. auth.
Suarez-Sipmann F. See *Tusman G,* jt. auth.
Sugino S, Kanaya N, Mizuuchi M, Nakayama M, Namiki A. Forehead is as sensitive as finger pulse oximetry during general anesthesia, 432
Sun W-Z. See *Lin C-S,* jt. auth.
Supandji M, Sia ATH, Ocampo CE. 0.2% ropivacaine and levobupivacaine provide equally effective epidural labour analgesia, 918
Suter PM. See *Bendjelid K,* jt. auth.

T
Takagi T. See *Iijima T,* jt. auth.
Takasu H. See *Sasano N,* jt. auth.
Takeda S, Masuda R, Kanazawa T, Tomaru T. Esmolol attenuates hepatic blood flow responses during sodium nitroprusside-induced hypotension in dogs, 348
Tallon J. See *Kovacs G,* jt. auth.
Tamanai-Shacoori Z, Shacoori V, Vo Van J-M, Robert J-C, Bonnaure-Mallet M. Sufentanil modifies the antibacterial activity of bupivacaine and ropivacaine, 911
Taniguchi H. See *Ogata J,* jt. auth.
Tao P-L. See *Chow L-H,* jt. auth.
Taylor P. See *Duflo F,* jt. auth.
Teng Y-H. See *Ting C-K,* jt. auth.
Tessler MJ, Rochon AG, Shrier I. Desflurane does not accelerate recovery from operations of short duration: a practice audit, 222
Tétrault J-P. See *Cardinal V,* jt. auth.
Tetsu S. See *Sasano N,* jt. auth.
Tetzlaff J. See *Dilger JA,* jt. auth.
Teunkens A. See *Missant C,* jt. auth.
Ting C-K, Hu JS, Teng Y-H, Chang Y-Y, Tsou M-Y, Tsai S-K. Desflurane accelerates patient response during the wake-up test for scoliosis surgery, 393
Tinmouth A. See *Fergusson D,* jt. auth.
To EWH. See *Ho AMH,* jt. auth.
Tokinaga Y. See *Yu J,* jt. auth.
Tomaru T. See *Takeda S,* jt. auth.
Tonelli M. See *Rapchuk I,* jt. auth.
Tong D. See *Chung F,* jt. auth.
Topcuoglu MS. See *Gulev T,* jt. auth.
Tourreau-Pham S. See *Fattorutto M,* jt. auth.
Townsend DR. See *Skjodt NM,* jt. auth.
Trachsel D. See *Dillier CM,* jt. auth.
Trager G. See *Hemmerling TM,* jt. auth.
Trenchant A. See *Van der Linden PJ,* jt. auth.
Tsai S-K. See *Ting C-K,* jt. auth.
Tsou M-Y. See *Ting C-K,* jt. auth.
Tsueda K. See *Chua MV,* jt. auth.
Tsui BCH, Wagner A, Finucane B. The threshold current in the intrathecal space to elicit motor response is lower and does not overlap that in the epidural space: a porcine model, 690
Turchetto E. See *Tusman G,* jt. auth.
Turcotte C, Drolet P, Girard M. Study design, originality and overall consistency influence acceptance or rejection of manuscripts submitted to the Journal, 549
Tusman G, Böhm SH, Suarez-Sipmann F, Turchetto E. Alveolar recruitment improves ventilatory efficiency of the lungs during anesthesia, 723

U
Unlugenc H. See *Gulev T,* jt. auth.

V
Vallejo MC. See *Ray BK,* jt. auth.
Van de Velde M. See *Missant C,* jt. auth.
Van der Linden PJ, De Hert SG, Daper A, Trenchant A, Schmartz D, Defrance P, Kimbimbi P. 3.5% urea-linked gelatin is as effective as 6% HES 200/0.5 for volume management in cardiac surgery patients, 236
van Eijs F. See *Castagnini HE,* jt. auth.
van Vlymen J. See *Bryson GL,* jt. auth.
VanDenKerkhof EG, Parlow JL, Goldstein DH, Milne B. In Canada, anesthesiologists are less likely to respond to an electronic, compared to a paper questionnaire, 449
VanDenKerkhof EG. See *Goldstein DH,* jt. auth.
Vandermeersch E. See *Missant C,* jt. auth.
Vandroux D. See *Nathan N,* jt. auth.
Varin F. See *Kaloul I,* jt. auth.
Vartazarian TN. See *Hagberg CA,* jt. auth.
Vavilala MS. See *Lee LA,* jt. auth.
Vergnes C. See *Lopez S,* jt. auth.
Veytsman AM. See *Yao L,* jt. auth.
Visco E. See *Lee LA,* jt. auth.
Vo Van J-M. See *Tamanai-Shacoori Z,* jt. auth.
von Goedeck A. See *Keller C,* jt. auth.

W
Wagner A. See *Tsui BCH,* jt. auth.
Walter SD. See *Sinuff T,* jt. auth.
Wang L. See *Hall SRR,* jt. auth.
Wang X. See *Zhang X,* jt. auth.
Watson J. See *Dugas G,* jt. auth.
Watson NC. See *Maltby JR,* jt. auth.
Waxler B, Mondragon SA, Patel SN, Nedumgottil K. Intrathecal lidocaine and sufentanil shorten postoperative recovery after outpatient rectal surgery, 680
—. Prophylactic ondansetron does not reduce the incidence of itching induced by intrathecal sufentanil, 685
Weiss M. See *Dillier CM,* jt. auth.*; Dullenkopf A,* jt. auth.
Wermelt J. See *Biedler A,* jt. auth.
Wilhelm W. See *Biedler A,* jt. auth.
Williams S. See *Arcand G,* jt. auth.
Williams SR, Denault AY, Pellerin M, Martineau R. Vasopressin for treatment of shock following aprotinin administration, 169
Wong AK. Full scale computer simulators in anesthesia training and evaluation, 455
Wong AYC, Chan RSN, Irwin MG. Anesthetic management of Cesarean delivery in a patient with hypoplastic anemia and severe pre-eclampsia, 923
Wong CL. See *Schwarz SKW,* jt. auth.
Wong DT. See *Friedman Z,* jt. auth.

X
Xiong L. See *Zhang X,* jt. auth.

Y
Yang H, Choi PT-L, McChesney J, Buckley N. Induction with sevoflurane-remifentanil is comparable to propofol-fentanyl-rocuronium in PONV after laparoscopic surgery, 660
Yao L, Veytsman AM, Dhamee MS. A right atrial foreign body (Images in anesthesia), 173
Yeh H-M. See *Lin C-S,* jt. auth.
Yoon S-M. See *Shin H-W,* jt. auth.
Yu J, Ogawa K, Tokinaga Y, Iwahashi S, Hatano Y. The vascular relaxing effects of sevoflurane and isoflurane are more important in hypertensive than in normotensive rats, 979
Yüceyar L, Erolçay H, Konukoglu D, Bozkurt K, Aykaç B. Epidural anesthesia may attenuate lipid peroxidation during aorto-femoral surgery, 465

Z
Zhang J. See *Munir MA,* jt. auth.
Zhang X, Xiong L, Hu W, Zheng Y, Zhu Z, Liu Y, Chen S, Wang X. Preconditioning with prolonged oxygen exposure induces ischemic tolerance in the brain via oxygen free radical formation, 258
Zheng Y. See *Zhang X,* jt. auth.
Zhu Z. See *Zhang X, jt.* auth.
Zschiegner F. See *Hinteregger M,* jt. auth.